Table 68. Healthy weight, overweight, and obesity among persons 20 years of age and over, according to sex, age, race, and Hispanic origin: United States, 1960–62, 1971–74, 1976–80, and 1988–94

[Data are based on measured height and weight of a sample of the civilian noninstitutionalized population]

Sex, age, race, and Hispanic origin ¹	Healthy weight ²				Overweight ³				Obesity ^A			
	1960–62	1971–74	1976–80 ⁵	1988–94	1960–62	1971–74	1976–80 ⁵	1988–94	1960–62	1971–74	1976–80 ⁵	1988–94
20–74 years, age adjusted ⁶	Percent of population											
Both sexes ^{7,8}		48.2	49.1	41.7	43.5	46.0	46.0	54.6	12.8	14.1	14.5	22.6
Male	48.1 52.1	44.0 52.2	46.0 52.1	39.1 44.3	48.4 38.6	52.7 39.7	51.3 40.8	59.4 49.9	10.4 14.9	11.8 16.1	12.2 16.5	19.9 25.1
White male	47.3 54.5	43.5 54.3	45.2 54.3	38.0 46.2	49.1 36.1	53.4 37.6	52.3 38.3	60.5 48.0	10.1 13.6	11.5 14.7	12.0 14.9	20.3 23.5
Black male	53.6 34.3	47.4 34.4	48.4 34.0	40.2 28.9	42.7 56.9	49.1 57.6	48.4 60.6	57.0 66.6	13.6 24.8	15.9 28.6	15.1 30.2	20.9 37.6
White, non-Hispanic male White, non-Hispanic female			45.5 54.9	38.7 48.0			52.0 37.6	59.9 45.7			11.9 14.8	20.1 22.5
Black, non-Hispanic male Black, non-Hispanic female ⁷			48.5 34.4	40.0 29.2			48.3 60.2	57.2 66.8			15.0 29.9	21.1 37.7
Mexican male			38.3 37.0	31.6 29.8			59.6 60.1	67.0 67.8			15.5 25.4	23.1 34.6
20-74 years, crude												
Both sexes ^{7,8}	49.0	47.5	48.7	41.4	45.2	47.0	46.4	55.0	13.5	14.4	14.7	22.7
Male	47.1 50.7	43.3 51.3	45.8 51.5	38.8 43.9	49.4 41.2	53.5 41.0	51.5 41.6	59.6 50.5	10.7 16.1	12.0 16.7	12.3 16.8	19.9 25.5
White male	46.3 53.0	42.6 53.3	45.0 53.6	37.5 45.4	50.2 38.9	54.3 39.1	52.5 39.4	61.1 49.0	10.4 14.7	11.7 15.4	12.1 15.3	20.4 24.0
Black male	53.0 33.2	47.3 34.2	48.3 34.5	40.6 29.7	43.9 58.8	49.3 58.2	48.5 60.0	56.7 65.9	14.1 26.6	16.0 28.7	15.0 29.8	20.9 37.0
White, non-Hispanic male \dots White, non-Hispanic female \dots			45.3 54.0	37.9 46.9			52.2 38.9	60.8 47.1			12.0 15.2	20.3 23.1
Black, non-Hispanic male Black, non-Hispanic female ⁷			48.4 35.0	40.4 29.9			48.4 59.4	57.0 66.2			14.9 29.5	21.1 37.2
Mexican male			41.0 39.4	34.4 31.2			57.0 57.4	64.0 66.2			14.6 23.8	20.7 33.6
Male												
20–34 years	44.1	53.9 34.5 36.7 38.1 41.4	55.7 40.5 37.9 37.9 41.1	50.3 33.3 33.5 28.1 29.8 40.6	42.7 53.5 53.9 52.2 47.8	42.8 63.2 59.7 58.5 54.6	41.2 57.2 60.2 60.2 54.2	47.5 65.5 66.1 70.5 68.5 56.5	9.2 12.1 12.5 9.2 10.4	9.7 13.5 13.7 14.1 10.9	8.9 13.5 16.7 14.1 13.2	14.1 21.5 23.2 27.2 24.1 13.2
Female ⁷												
20–34 years		61.7 53.5 47.7 39.2 39.6	61.0 53.4 47.2 42.7 36.5	54.3 45.5 35.6 31.2 36.0 41.0	21.2 37.2 49.3 59.9 60.9	25.8 40.5 49.0 54.5 55.9	27.9 40.7 48.7 53.7 59.5	37.0 49.6 60.3 66.3 60.3 52.3	7.2 14.7 20.3 24.4 23.2	9.7 17.7 18.9 24.1 22.0	11.0 17.8 19.6 22.9 21.5	18.5 25.5 32.4 33.7 26.9 19.2

^{- - -} Data not available.

NOTES: Percents do not sum to 100 because the percent of persons with BMI less than 19 is not shown and the percent of persons with obesity is a subset of the percent with overweight. Height was measured without shoes; two pounds are deducted from data for 1960–62 to allow for weight of clothing.

SOURCE: Centers for Disease Control and Prevention, National Center for Health Statistics, Division of Health Examination Statistics. Unpublished data.

¹The race groups, white and black, include persons of Hispanic and non-Hispanic origin.

²Body mass index (BMI) of 19 to less than 25 kilograms/meter² (see Appendix II, Body mass index).

³BMI greater than or equal to 25.

⁴BMI greater than or equal to 30.

⁵Data for Mexicans are for 1982–84. See Appendix I.

⁶See Appendix II for age-adjustment procedure.

⁷Excludes pregnant women.

⁸Includes persons of all races and Hispanic origins, not just those shown separately.